

<i>Anti-gas</i>							
DRUGS	TRADE NAME	ACTIVE INGREDIENT	INDICATIONS	DOSAGE/CONCENTRATION & ROUTES OF ADMIN.	TOXIC/LETHAL DOSE	WITHDRAWAL TIME	CONTRADICTIONS
Ventipulmin	Ventipulmin® Syrup	Clenbuterol Hydrochloride	Management of horses affected with airway obstruction, eg: chronic obstructive pulmonary disease (COPD).	Orally twice daily with initial dosage at 0.5 mL/100 lbs of body weight (0.8 mcg/kg)	Oral (rat) LD50: 159 mg/kg	72 hours	antagonizes the effects of prostaglandin F2 alpha and oxytocin. Ventipulmin Syrup should not be used in pregnant mares near term. Because tachycardia may occur, Ventipulmin Syrup should not be used in horses suspected of having cardiovascular impairment.
Scourban Plus	Scourban®	Sulphadimidine 639mg, Sulphaguanidine 639mg, Sulphadiazine 852mg, Hyoscine	Prevention and treatment of intestinal infections of bacterial origin in horses, cattle, goats, pigs, sheep, dogs and cats.	For oral use only. Small animals: 4 – 8mL per 8 – 10kg bodyweight Calves, Pigs & other large animals: 30mL per 25kg bodyweight Piglets: 1 – 2 mL Lambs: 5mL		MEAT: 14 days. MILK: 35 days.	Not for use in bobby calves.
PCE-Glycol		Propylene Glycol 0.667 mL Choline Chloride 20.00 mg Potassium Iodide 1.76 mg Ethylenediamine Dihydroiodide 0.44 mg Cobalt Sulfate	Glucogenic as an aid in the prevention and treatment of acetonemia	Prevention Cattle 175 - 225 mL daily. Begin 2 weeks prior to calving and continue for 6 weeks after calving. Sheep 50 - 100 mL daily. Begin 4 weeks prior to lambing and continue until lambing. Treatment Cattle 200 - 400 mL daily, dependant upon severity. Administer for 10 days. Sheep 75 - 125 mL daily, dependant upon severity. Administer for 10 days.		-	