

DRUG	ACTIVE INGREDIENT	USE	ROUTE OF ADMINISTRATION	ANIMAL TREATED	DOSAGE	WARNINGS	WITHDRAWAL PERIOD
Progestrin REP	progesterone	Given to female animals To increase endometrial receptivity for implantation of an embryo	Intra muscularly Orally	Horse Cattle Small ruminants Pig	Progesterone USP, in oil should be given intramuscularly daily as follows: Cows and Mares - 500 mg Sheep and Swine – 250 mg		
Oxyto-Kel 10 IU	Oxytocin 10 IU	to increase uterine contractions and/or milk letdown.	Intra muscular Intravenous	Horse Cattle Small ruminants Pig	In horse via IM: 5ml /500kg Body Weight (B.W.) Via IV: 1-2 ml/500 kg B.W. In Cow Via IM:2ml /500kg B.W. Via IV: 1ml/500 B.W	This product must not be used in lactating dairy animals.	ZERO days

					In Goat: 1 ml/50 kg B.W In Pig: Via Im:1-2 ml /200kg B.w. I.V .5ml/200kg / B.W.		
ECP 2mg/ml	Estradiol cypionate	The active ingredient activates the transcription of genes involved in the functioning of the female reproductive system and secondary sex characteristics.	Intra muscular	Horse Cows	In horse 5-10mg/ ml (as stated in the picture) . In cows 2mg/ml ((as stated in the picture)	Injection site reactions may occur. Precautions Estrogens generally are contraindicated in pregnant mares.	ZERO days
Lutalyse	prostaglandin F2 alpha (dinoprost) as the tromethamine salt.	For intramuscular use for estrus synchronization, treatment of unobserved (silent) estrus and pyometra (chronic endometritis) in cattle; for abortion of	Intra muscular	Beef Cattle Dairy Cattle Horses Swine	In horse via IM: Via IV In Cow Via IM for estrus synchronization: Inject a dose of 5	Residue Warnings: . Use of this product in excess of the approved dose may result in drug residues. Do not use in horses	No milk discard or preslaughter drug withdrawal period is required for labeled uses in cattle or Swine .

		feedlot and other non-lactating cattle; for parturition induction in swine; and for controlling the timing of estrus in estrous cycling mares and clinically anestrous mares that have a corpus luteum.			mL LUTALYSE (25 mg PGF2 α) either once or twice at a 10 to 12 day interval For Intramuscu lar Use for Treatment of Pyometra (chronic endometrit is) in Cattle. Inject a dose of 5 mL LUTALYSE (25 mg PGF2 α) For Intramuscu lar Use for Abortion of Feedlot and Other Non- Lactating Cattle.:	intended for human consumption.	
--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------	--

					<p>Inject a dose of 25 mg intramuscularly.</p> <p>In Goat In Pig: For intramuscular use for parturition induction in swine. LUTALYSE Sterile Solution is indicated for parturition induction in swine when injected within 3 days of normal predicted farrowing.</p>	
Fertiline solution	gonadorelin acetate	Used in bovine estrous synchronization program	Intra muscular	Cattle	<p>In Cow Via IM: 100 µg per cow (2 mL)</p>	Zero milk withdrawal

					of a solution containing 50 µg/mL).		
Metricyclin [chlortetracycline HCL 1g]	Chlortetracycline hydrochloride 1 g	For the prevention of endometritis after parturition and for the treatment of uterine infections caused by germs sensible to chlortetracycline	bolus for intrauterine use	Cows, ewes	For intra-uterine administration shortly after calving. Prophylactic: 1 bolus Therapeutic: 1-2 boluses	Dangerous to the Environment.	Milk: Milk intended for sale for human consumption must be discarded during treatment and for not less than 2 milkings or approximately 24 hours following the last treatment. Meat: Animals Producing Meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.
Metricure 500mg [intrauterine suspension 19G]	cefapirin (as cefapirin benzathine)	for the treatment of subacute and chronic endometritis in cows (at least 14 days after parturition) caused by bacteria sensitive	intrauterine	Cattle	The contents of one Metricure syringe should be introduced into the lumen of	Not to be administered to animals known to be allergic to cephalosporins or penicillins. The product is not indicated	Meat - 1 day after last treatment. Milk - Zero hours.

		to cefapirin.			the uterus using a catheter. One treatment with Metricure is usually sufficient for a complete cure. In animals that have been inseminated, Metricure may be used at one day after insemination	for use during pregnancy but can be used during lactation. Not to be used concurrently with other intra-uterine antibiotic preparations.	
Eazi-Breed CIDR	Progesterone,	Uses: method for both synchronising and inducing oestrus in production animals. ● Synchronization of estrus in lactating dairy cows, suckled	Intravaginally placed	Cattle Sheep Goats	Goats: Insert Eazi-Breed CIDR devices during the breeding season and leave in place for 18 to 21	Store in a dry place below 30°C (room temperature). void contact with skin by wearing protective gloves when handling the	For sheep and goats: Withholding period: * Meat: 24 hours

		<p>beef cows, and replacement beef and dairy heifers</p> <ul style="list-style-type: none"> ● Induction of estrous cycles in anestrous lactating dairy cows ● Synchronization of the return to estrus in lactating dairy cows inseminated at the immediately preceding estrus ● Advancement of first postpartum estrus in suckled beef cows ● Advancement of first pubertal estrus in replacement beef heifers 			<p>days.</p> <p>Cattle :1.38 grams per EAZI-BREED CIDR Cattle Insert</p> <p>Sheep: Insert Eazi-Breed CIDR devices during the breeding season and leave in place for 12 to 14 days.</p>	<p>inserts. Not for use in humans. Keep out of reach of children. Store used (removed) EAZI-BREED CIDR Cattle Inserts in a plastic bag or other sealable container until they can be properly disposed in accordance with applicable local, state and Federal regulations.</p>	
--	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--